

RL-2582 (6/08)

EXP.

LOT

MFD



LIFT  
HERE

50 mL Single patient infusion vial

**PROPOFOL**  
Injectable Emulsion  
500 mg/50 mL  
(10 mg/mL) **1%**

**CONTAINS BENZYL ALCOHOL  
FOR I.V. ADMINISTRATION  
SHAKE WELL BEFORE USING.**

Hospira, Inc., Lake Forest, IL 60045 USA

- **CONTAINS BENZYL ALCOHOL, which inhibits microbial growth up to 12 hours**
- **Use strict aseptic technique**
- **Contamination can cause fever, infection/sepsis and/or other life-threatening illness**
- **Begin use promptly after opening; Discard within specified time limit (see package insert)**
- **Do not use if contamination is suspected**

Printed in USA

NDC 0409-4699-33

Each mL contains propofol 10 mg; soybean oil 100 mg; glycerin 22.5 mg; egg lecithin 12 mg, benzyl alcohol 1.5 mg and sodium benzoate 0.7 mg added. pH adjusted with sodium hydroxide. Sterile, nonpyrogenic. **USUAL DOSAGE:** See insert. Store at 20 to 25°C (68 to 77°F). [See USP Controlled Room Temperature.] Do not freeze. Discard unused portion.

Rx only

  
**Hospira**