

signs of methemoglobinemia, a rare disorder, which may appear up to 2 hours after use.

• **Do not exceed recommended dosage.**

If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center immediately.

• **If pregnant or breast feeding,** ask a health professional before use.

• **Keep out of reach of children.**

Directions

Apply only amount needed to the oral mucosa to prevent or relieve pain.

Other information

Store at 59° - 86°F (15 - 30°C). Protect from freezing.

Inactive ingredients

flavoring, PEG 3350, PEG 400, sodium saccharine. May contain yellow #5 (tartrazine), yellow #6, red #3, red #40, blue #1, green #3 as a color additive.

Questions or comments?

800.873.7683

Manufactured for:

Fabrique pour :

Patterson Dental Supply, Inc.
1031 Mendota Heights Road
Saint Paul, MN 55120

Drug Facts

**Active ingredients
(in each gram)**

Benzocaine 200 mg

Purpose

Oral Anesthetic

Use

For oral mucosal use only, as directed by dentist. For the temporary relief of pain due to minor dental procedures.

Warnings

- **Allergy Alert:** Do not use on patients with a history of allergies to local anesthetics such as benzocaine or other "caine" anesthetics.
- **Do not use** for more than 7 days unless directed by a dentist or doctor. If sore mouth symptoms do not improve in 7 days; irritation, pain, or redness persists or worsens; or if swelling, rash or fever develops, see your dentist or doctor promptly.
- **When using this product** Avoid contact with eyes. If contact occurs, flush with water.
- **Stop use and consult a health care practitioner** if the following symptoms appear: weakness, confusion, headache, difficulty breathing, and/or pale, grey or blue colored skin, as these may be

P151153 (1/15)

**Topical
Anesthetic Gel**
1 oz (30 ml)


PATTERSON
DENTAL

Manufactured for:
Fabrique pour :

PATTERSON
DENTAL
Patterson Dental Supply, Inc.
1031 Mendota Heights Road
Saint Paul, MN 55120