



DISCOVER Orabloc®

(Articaine HCl 4% and epinephrine 1:100,000 and epinephrine 1:200,000) injection


The Quality that you need the Price that you want

“Articaine local anesthetic was listed as a top product dentists couldn't live without. Presently, Orabloc by Pierrel Pharma has lower cost and longer shelf-life” *

*An independent, non-profit, dental education and product testing foundation. Clinicians Report, December, 2014. For the full report go to www.Orabloc.com



The Lowest Price and the Longest Shelf-life



- Rapid onset of anesthesia within 1-9 minutes
- Complete anesthesia lasts about 1 hour for infiltration, up to 2 hours for nerve block
- Only 10% overage of epinephrine¹
- 24 month shelf life at room temperature
- Sodium edetate free, methylparaben free and latex free
- Each cartridge is sealed individually in the blister for maximum protection

¹The American Heart Association (AHA) recommends using the lowest possible quantity of epinephrine (Kaplan EL ed. Cardiovascular disease in dental practice. Dallas, TX: American Heart Association, 1986)

(articaine HCl 4% and epinephrine 1:100,000 and epinephrine 1:200,000) Injection

Orabloc® is indicated for local, infiltrative, or conductive anesthesia in both simple and complex dental procedures:

- For most routine dental procedures, Orabloc® containing epinephrine 1:200,000 is preferred
- When more pronounced homeostasis or improved visualization of the surgical field are required, Orabloc® containing epinephrine 1:100,000 may be used

Dosage and administration – Adults

- For normal healthy adults, the maximum dose of Orabloc® administered by submucosal infiltration and/or nerve block should not exceed 7mg/kg (0.175 mL/kg) of articaine HCl
- Dosage should be reduced in elderly patients and in patients with cardiac or liver disease

Pediatric patients ages 4 to 16 years

- The quantity of Orabloc® in children ages 4 to 16 years of age to be injected should be determined by the age and weight of the child and the magnitude of the operation
- The maximum dose of Orabloc® should not exceed 7 mg/kg (0.175 mL/kg) of articaine HCl (see Use in Specific Populations). Use in pediatric patients under 4 years of age is not recommended

Both Orabloc® strengths have a 24 month shelf life

- Store at room temperature; 25°C (77°F), with brief excursions permitted between 15°C (59°F) and 30°C (86°F)
- Protect from light
- Do not freeze

Orabloc® packaging

- Each cartridge is individually sealed for maximum protection up to the moment of use
- Cartridges packed 10 to a blister tray to avoid glass to glass contact
- Blister trays packaged in boxes of 50

Administration of Orabloc results in a 3 to 5 fold increase in plasma epinephrine concentrations compared to baseline.

However, in healthy adults it does not appear to be associated with marked increases in blood pressure or heart rate, except in the case of accidental intravascular injection.

Inform patients in advance of the possibility of temporary loss of sensation and muscle function following infiltration and nerve block injections. Instruct patients not to eat or drink until normal sensation returns.

To know where to order link to www.orabloc.com/contact-us.html
To learn more about Orabloc®, call Pierrel at 1-610-989-4222

